

## QUARKONIUM AT FINITE TEMPERATURE \*


P. PETRECZKY <sup>†</sup>

*Physics Department, Brookhaven National Laboratory, Upton, NY 11973, USA*

I discuss quarkonium spectral functions at finite temperature reconstructed using the Maximum Entropy Method. I show in particular that the  $J/\psi$  survives in the deconfined phase up to  $1.5T_c$

The study of quarkonium system at finite temperature has been a subject of considerable interest since the work of Matsui and Satz <sup>1</sup>, but a first principle calculation of quarkonium properties at non-zero temperature was missing. It was shown, however, that the application of the Maximum Entropy Method (MEM) can make such calculation possible <sup>2</sup>. The method have been successfully applied at zero <sup>2</sup> as well as at finite temperature <sup>3</sup>.

I am going to discuss charmonium spectral function calculated with MEM on  $48^3 \times N_\tau$  lattices at lattice spacing  $a^{-1} = 4.86\text{GeV}$  and  $N_\tau = 24, 16$  and  $12$  corresponding to temperatures  $0.75T_c$ ,  $1.12T_c$  and  $1.5T_c$  ( $T_c$  being the deconfinement temperature). The results for the vector channel are shown in the Figure. As one can see the  $J/\psi$  seems to survive up to temperatures  $1.5T_c$ . Similar calculation have been performed in the scalar and axial vector channels which correspond to the  $P$ -state charmonia, but no peak was found there.



### References

1. T. Matsui and H. Satz, *Phys. Lett.* **B178**, 416 (1986).
2. M. Asakawa et al., *Prog. Part. Nucl. Phys.* **46**, 459 (2001).
3. F. Karsch et al., *Phys. Lett.* **B530**, 147 (2002); S. Datta et al., hep-lat/0208012; M. Asakawa and T. Hatsuda, hep-lat/0308034.

\*Based on work done in collaboration with S. Datta, F. Karsch and I. Wetzorke.

<sup>†</sup>Goldhaber fellow, supported under contract DE-AC02-98CH10886 with the U.S. Department of Energy.